

Chad J. Muxlow, D.O.

Director of Sports Medicine Surgery,
Sentara RMH Orthopedics and Sports Medicine
Head Team Surgeon, Bridgewater College
Head Team Surgeon, Eastern Mennonite University
www.MuxlowSportsMedicine.com



MUXLOW SPORTS MEDICINE

Physical Therapy Prescription – SLAP Repair

MOON (Multicenter Orthopaedic Outcomes Network) Protocol

Name: _____ **Date:** _____

Diagnosis: R / L arthroscopic SLAP repair **Date of Surgery:** _____

Frequency: 2-3 times per week for _____ weeks, **beginning 1 week after surgery**

Phase I (Weeks 0 – 6):

- **Sling with abduction pillow:** Continue for a total of 6 weeks
- **Range of Motion:**
 - Weeks 1-2: PROM, AAROM including FF to 60°, ER to neutral with arm at side, IR to 45°
 - i. **NO active ER, extension, or abduction**
 - Weeks 3-4: PROM, AAROM including FF to 90°, ABD to 85°, ER at 30° of ABD to 30°, IR at 30° of ABD to 60°
 - i. **NO active ER or extension**
 - Weeks 5-6: PROM, AAROM including FF to 145°, ER at 45° of ABD to 50°, IR at 45° of ABD to 60°
 - Week 6: initiate gentle ROM at 90° of abduction, progress to 30° of ER
- **Exercises:** begin gentle isometrics at week 2; but no ER/IR
 - Weeks 3-4: begin scapular stabilizers (protraction, retraction) with arm in sling
 - Weeks 5-6: initiate active shoulder ABD (without resistance), “full can” exercises, prone rowing, prone horizontal abduction
 - Week 6: start biceps isotonic
- **Modalities:** Per therapist, including electrical stimulation, ultrasound, heat (before), ice (after)

Phase II (Weeks 7 – 14):

- **Range of Motion:** increase as tolerated, goal is full ROM by week 10, full AROM by week 12
 - Week 10-11: Progress ROM to functional demands (ie, overhead athlete, throwing motions)
- **Exercises:** continue Phase I; progress isotonic strengthening program, PNF strengthening, stretching
 - Weeks 7-9: Initiate Throwers Ten Program
- **Modalities:** Per therapist, including electrical stimulation, ultrasound, heat (before), ice (after)

Phase III (Weeks 15 – 20):

- **Exercises:** continue Phase II, advance as tolerated
 - Weeks 14-16: initiate light plyometric program, restricted sports activities including light swimming and half-golf swings
 - Week 16: initiate interval sport program (ie, throwing)
- Okay to return to contact sports or heavy labor at 20 weeks if patient has full non-painful ROM, satisfactory static stability, muscular strength 75-80% contralateral side, and no pain/tenderness

Phase IV (Weeks 20+):

- **Exercises:** continue Phase III, advance as tolerated, progress interval sport programs
- Okay to return to pitching if successful completes interval throwing program without pain, and has full functional ROM, satisfactory stability, and no pain/tenderness.

Signature: _____

Date: _____